arXiv:1201.4635v1 [nucl-ex] 23 Jan 2012

Kaonic hydrogen X-ray measurement in SIDDHARTA

M. Bazzi[a], G. Beer[b], L. Bombelli[c], A.M. Bragadireanu[a,d], M. Cargnelli[e], G. Corradi[a],
C. Curceanu (Petrascu)[a], A. d'Uffizi[a], C. Fiorini[c], T. Frizzi[c], F. Ghio[f], C. Guaraldo[a],
R.S. Hayano[g], M. Iliescu[a,d], T. Ishiwatari[e], M. Iwasaki[h], P. Kienle[e,i], P. Levi Sandri[a],
A. Longoni[c], V. Lucherini[a], J. Marton[e], S. Okada[a,*], D. Pietreanu[a,d], T. Ponta[d], A. Rizzo[a],
A. Romero Vidal[a], A. Scordo[a], H. Shi[g], D.L. Sirghi[a,d], F. Sirghi[a,d], H. Tatsuno[g,1],
A. Tudorache[d], V. Tudorache[d], O. Vazquez Doce[a], E. Widmann[e], J. Zmeskal[e]

[a] *INFN, Laboratori Nazionali di Frascati, Frascati (Roma), Italy*
[b] *Department of Physics and Astronomy, University of Victoria, Victoria, BC, Canada*
[c] *Politecnico di Milano, Sezione di Elettronica, Milano, Italy*
[d] *Horia Hulubei National Institute of Physics and Nuclear Engineering, Magurele, Romania*
[e] *Stefan-Meyer-Institut für subatomare Physik, Vienna, Austria*
[f] *INFN Sezione di Roma I and Istituto Superiore di Sanità, Roma, Italy*
[g] *University of Tokyo, Tokyo, Japan*
[h] *RIKEN, The Institute of Physics and Chemical Research, Saitama, Japan*
[i] *Technische Universität München, Physik Department, Garching, Germany*

Abstract

Kaonic hydrogen atoms provide a unique laboratory to probe the kaon-nucleon strong interaction at the energy threshold, allowing an investigation of the interplay between spontaneous and explicit chiral symmetry breaking in low-energy QCD. The SIDDHARTA Collaboration has measured the K-series X rays of kaonic hydrogen atoms at the DAΦNE electron-positron collider of Laboratori Nazionali di Frascati, and has determined the most precise values of the strong-interaction induced shift and width of the $1s$ atomic energy level. This result provides vital constraints on the theoretical description of the low-energy $\overline{K}N$ interaction.

Keywords: Kaonic atoms, Low-energy QCD, Antikaon-nucleon physics, X-ray detection

1. Introduction

Kaonic hydrogen is an exotic atom composed of a proton and a K^- bound by the Coulomb force. The strong interaction shifts the $1s$ atomic energy level from its pure electromagnetic (EM) value and broadens it as a result of the shortened lifetime due to absorption. Since the strong-interaction effects for the higher states (*i.e.*, $2p$, $3p$ ...) are negligible in comparison, the $1s$ shift and width can be deduced from the spectroscopy of kaonic-hydrogen X-ray transitions feeding the $1s$ states, namely the K-series X rays.

---

[*] Corresponding author.
Email address: `shinji.okada@lnf.infn.it` (S. Okada)
[1] Present address: INFN, Laboratori Nazionali di Frascati, Frascati (Roma), Italy

Preprint submitted to Nuclear Physics A *January 24, 2012*

The measured strong-interaction shift ϵ_{1s} and width Γ_{1s} are directly related to the real and imaginary parts of the complex $K^- p$ S-wave scattering length a_{K^-p} which, in the isospin limit, is given by the Deser-Truemann formula [1] :

$$\epsilon_{1s} + \frac{i}{2}\Gamma_{1s} = 2\alpha^3\mu^2 a_{K^-p} = 412 \text{ eV fm}^{-1}\, a_{K^-p}$$

where μ is the reduced mass of the $K^- p$ system and α is the fine structure constant. The kaonic-hydrogen X-ray data are therefore crucial for theories of the $\overline{K}N$ system together with the low-energy $\overline{K}N$ data. Note that recent Coulomb and isospin breaking corrections to the formula turn out to be important, as first shown in [2]. Fig. 1 shows the comparison between uncorrected and corrected values in this theoretical calculation [2].

The low-energy $\overline{K}N$ system has attracted attention as a sensitive testing ground for chiral SU(3) dynamics in low-energy QCD, allowing investigation of the interplay between spontaneous and explicit chiral symmetry breaking due to the relatively large strange quark mass which plays an intermediate role between "light" and "heavy". The data are also strongly related to recent hot topics – the structure of the $\Lambda(1405)$ resonance (*e.g.*, [3, 4, 5]) and the deeply bound kaonic systems (*e.g.*, [6, 7, 8, 9]). Recent progress in this field is summarized in [10].

Historically there were several measurements of the strong-interaction shift ϵ_{1s} [2] and width Γ_{1s} [11, 12, 13, 14, 15]. In the 1970s and the 1980s three groups [11, 12, 13] reported a measured attractive shift (positive ϵ_{1s}) as shown in Fig. 1, while the information extracted from the analyses of the low energy $\overline{K}N$ data (*e.g.*, [16, 17, 18]) shows a repulsive shift (negative ϵ_{1s}). This contradiction has been known as the "kaonic hydrogen puzzle".

In 1997, the first distinct peaks of the kaonic-hydrogen X rays were observed by the KEK-PS E228 group [14] with a significant improvement in the signal-to-background ratio by the use of a gaseous hydrogen target, where previous experiments had employed liquid hydrogen. It was crucial to use a low-density target, namely a gaseous target, because the X-ray yields quickly decrease towards higher density due to the Stark mixing effect. The observed repulsive shift was consistent in sign with the analysis of the low energy $\overline{K}N$ scattering data, resolving the long-standing discrepancy.

More recent values reported by the DEAR group in 2005 [15], with substantially reduced errors as shown in Fig. 1, firmly established the repulsive shift obtained in the previous E228 experiment.

Intensive theoretical studies were performed based on these DEAR results [2, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28] [3]. However, most calculations show difficulties in explaining all the experimental results in a consistent way. As an example, the theoretical values with and without restriction of DEAR data are plotted in Fig. 1 [2].

In the present SIDDHARTA experiment we have determined the most precise values to date of the $1s$ strong interaction shift and width of kaonic hydrogen [31] in an attempt to clarify this difficulty.

2. Experiment

The SIDDHARTA experiment was performed at the recently upgraded DAΦNE positron-electron collider [32, 33]. The collider produces ϕ-resonances, 49 % of which decay into back-

---

[2] Note that ϵ_{1s} is defined as $\epsilon_{1s} \equiv -(E_{1s} - E^{EM}_{1s})$, where E_{1s} is the energy of the $1s$ level and E^{EM}_{1s} is the energy calculated using only the EM interaction.

[3] See also [29, 30].



Figure 1: Comparison of previous experimental and theoretical results for the strong-interaction 1*s*-energy-level shift and width of kaonic hydrogen. The open circles with error bars are experimental results: J. D. Davies *et al.* (1979) [11], M. Izycki *et al.* (1980) [12], P. M. Bird, *et al.* (1983) [13], KEK-PS E228 (1997) [14] and DEAR (2005) [15]. The error bars correspond to quadratically added statistical and systematic errors. The symbols without error bars show the values of a theoretical calculation [2] both by using the Deser-Trueman formula (empty symbols) and by including isospin breaking corrections (filled symbols). The fit restricted to the DEAR data is represented by the squares.

to-back K^+K^- pairs. The resulting monochromatic low-energy kaons are efficiently stopped in a cryogenic hydrogen "gaseous" target.

A schematic view of the SIDDHARTA setup is shown in Fig. 2. A coincidence of two plastic scintillation counters mounted above and below the e^+e^- interaction point was used as a kaon trigger. X rays emitted from the kaonic atoms were detected by 144 silicon drift detectors (SDDs), each having an effective area of 1 cm^2 and a thickness of 450 μm, developed within a European research project devoted to this experiment. The SDDs had an energy resolution of ~ 180 eV (FWHM) at 8 keV and timing resolution below 1 μsec, in contrast to the CCD detectors used in DEAR [34] which had no timing capability. In comparison with DEAR, the main source of background coming from beam losses was highly suppressed. A detailed description of our experimental setup is given in [31, 35, 36].

3. Data analysis

Fig. 3 shows the timing distribution of the coincidence signals in the kaon detector with respect to the ~ 368.7 MHz RF signal from DAΦNE. The spectrum shows clearly that kaon events can be separated from minimum ionizing particles by setting a time gate as indicated by arrows in the figure. A half frequency of the beam synchronous timing signal was used for the start RF timing – therefore there are two identical coincidence signals in the spectrum at an interval of ~ 2.6 ns.

Energy calibration of each SDD's ADC spectrum was performed by periodic measurements of characteristic X-ray lines from titanium and copper foils located as shown in the setup (Fig. 2) excited by an X-ray tube. Fig. 4 (a) shows a typical X-ray spectrum from the X-ray tube data for a single SDD. Characteristic X rays of titanium and copper were obtained with high statistics. A remote-controlled system moved the kaon detector out and the X-ray tube in for these calibration



Figure 2: A schematic view of the SIDDHARTA setup installed at the e^+e^- interaction point of DAΦNE.

measurements, once every ∼ 4 hours. The energy scale was calibrated linearly by the two $K\alpha$ lines.

The refined *in-situ* calibration in energy was performed using characteristic X-ray lines of titanium, copper and gold ($L\alpha$) excited by the uncorrelated background without trigger in the summed spectrum of all SDDs. The spectrum for the complete statistics of the kaonic-hydrogen dataset is shown in Fig. 4 (b). Moreover the parameters of the (energy-dependent) energy resolution for the summed spectrum were also evaluated using those peaks in this *in-situ* energy spectrum along with the kaonic carbon lines from wall stops in the final energy spectrum in the kaon-triggered mode.

A correlation plot of the X-ray energy measured by SDDs vs the time difference between kaon arrival (with kaon detectors) and X-ray detection (with SDDs) for hydrogen data is shown in Fig. 5. The horizontal band is due to the kaon-induced events. Its projected time and energy spectra are also displayed in the figure. A typical width of the time-correlation, after the time-walk correction, was about 800 ns (FWHM) which reflected the drift-time distribution of the



Figure 3: Kaon identification using timing of the coincidence signals in the kaon detector with respect to the RF signal of ~ 368.7 MHz from DAΦNE.

electrons in the SDD. The kaon and background time gates, each having a width of 1 μsec, are indicated therein with arrows.

The continuous background is related to the following two types of particles: the charged kaon secondaries (prompt background) and lost beam particles (accidental background). Comparing the energy spectra of data with SDD timing gates set to "K" (kaon) and "BG" (background) in Fig. 5, the prompt background is at the same level as the accidental background. In the most recent previous measurement of kaonic hydrogen X rays (DEAR) [15], also performed at DAΦNE, the kaonic-hydrogen spectrum suffered from the huge accidental background due to lack of the timing capability of the X-ray detectors (CCDs) used. The event selection using the time information significantly reduced the accidental background and improved the signal-to-background ratio by more than a factor of 10 with respect to the corresponding DEAR ratio of about 1/100.

Fig. 6 (b) shows the kaonic-hydrogen X-ray spectrum. We have also measured an X-ray spectrum with a deuterium target (for the first-ever exploratory measurement of kaonic-deuterium X rays), as shown in Fig. 6 (c). The kaonic-hydrogen X-ray transitions were clearly observed while those for kaonic deuterium were not visible. This appears to be consistent with the theoretical expectation that kaonic deuterium X rays have one order lower yield per stopped K^- and greater width than those of kaonic hydrogen X rays (*e.g.*, [37]).

A dot-dashed line in Fig. 6 (b) indicates the EM value of the kaonic-hydrogen $K\alpha$. Comparing the kaonic-hydrogen $K\alpha$ peak and the EM value, there is no room for doubt about a repulsive shift of the kaonic-hydrogen 1s-energy level, which is consistent with the analysis of the low energy $\overline{K}N$ scattering data.

Many other kaonic-atom X rays and characteristic X rays were detected in both spectra as indicated with arrows in the figures. Those kaonic-atom lines are attributable to the target-cell wall made of Kapton polyimide film ($C_{22}H_{10}O_5N_2$) and its support frames made of aluminum. The characteristic X rays come from high-purity titanium and copper foils installed for *in-situ* X-ray energy calibration.

There are three background X-ray lines overlapping with the kaonic-hydrogen signals : kaonic oxygen 7-6 (6.0 keV), kaonic nitrogen 6-5 (7.6 keV) and the characteristic X ray of



Figure 4: X-ray energy spectra for (a) data taken with the X-ray tube, and (b) data uncorrelated to the kaon production timing. The refined *in-situ* calibration in gain (energy) and resolution (response shape) of the summed spectrum of all SDDs was obtained using titanium, copper, and gold characteristic X-ray lines in the spectrum (b).

copper $K\alpha$ (8.0 keV). In the fitting procedure of the kaonic-hydrogen spectrum, it turned out to be essential to use the kaonic-deuterium spectrum to quantify the kaonic background X-ray lines. Therefore, we performed a simultaneous global fit of the hydrogen and deuterium spectra, where the intensities of the background X-ray lines were determined using both spectra and a normalization factor defined by the intensity ratio of the high-statistics kaonic-carbon 5-4 peak seen in both spectra. In Fig. 6 (b) and (c), the resulting fit lines are shown together with components of both the background X-ray lines and a continuous background; (a) shows the residuals of the measured kaonic-hydrogen X-ray spectrum after subtraction of the fitted background, clearly displaying the kaonic-hydrogen K-series transitions.

As a result, the $1s$-level shift ϵ_{1s} and width Γ_{1s} of kaonic hydrogen were determined by SIDDHARTA to be

$$\begin{aligned} \epsilon_{1s} &= -283 \pm 36(\text{stat}) \pm 6(\text{syst})\ \text{eV} \\ \text{and}\ \ \Gamma_{1s} &= 541 \pm 89(\text{stat}) \pm 22(\text{syst})\ \text{eV}, \end{aligned}$$

respectively, where the first error is statistical and the second is systematic. The quoted systematic error is a quadratic summation of the following contributions : the SDD gain shift, the SDD response function, the ADC linearity, the low-energy tail of the kaonic-hydrogen higher transitions, the energy resolution, and the procedural dependence shown by an independent analysis [31].



Figure 5: Measured X-ray time and energy spectra for K^- triggered events of hydrogen data. The top-left panel shows a correlation plot of the X-ray energy measured by SDDs vs the time difference between kaon arrival and X-ray detection, where a time-walk correction was applied. The projections on the time and energy axes are also shown at the right and bottom. By selecting kaon and background time gates indicated with “K” and “BG” in the timing spectrum, both the energy spectra are displayed.

4. Conclusion

We have determined the strong-interaction energy-level shift and width of the kaonic-hydrogen atom 1*s* state with the best accuracy up to now [31]. The obtained shift and width are plotted in Fig. 7 along with the other two recent results [14, 15]. It should be noted that the smaller the width, the better the accuracy of determining the energy. The right panel of Fig. 7 shows the errors on the energy shift as a function of the width (vertical axis) for each experiment, together with guide lines representing SIDDHARTA precision calculated assuming the same statistics but with differing width. In comparison with the DEAR result, the accuracy of determining the energy in SIDDHARTA is obviously improved.

The new triggerable X-ray detectors, SDDs, developed in the framework of the SIDDHARTA project, lead to an improved energy and time resolution over the past experiments, resulting in much lower background in comparison with the DEAR experiment, thus permitting a better control on systematics.

Our determination of the shift and width allows more precise evaluation of $\overline{K}N$ scattering lengths which yields vital constraints on the theoretical description of the low-energy $\overline{K}N$ interaction (*e.g.*, [38, 39]).

For further study of the "isospin-dependent" $\overline{K}N$ interaction, it is crucial to measure the strong-interaction 1*s*-energy-level shift and width of kaonic deuterium. The present kaonic-hydrogen result combined with kaonic-deuterium data to be collected in the SIDDHARTA-2 experiment [40], being designed with more than one order-of-magnitude better signal-to-background ratio than that in SIDDHARTA, will provide invaluable constraints for the theories of low-energy QCD in the strangeness sector.

Acknowledgments

We thank C. Capoccia, B. Dulach, and D. Tagnani from LNF-INFN; and H. Schneider, L. Stohwasser, and D. Stückler from Stefan-Meyer-Institut, for their fundamental contribution in designing and building the SIDDHARTA setup. We thank as well the DAΦNE staff for the excellent working conditions and permanent support. Part of this work was supported by HadronPhysics I3 FP6 European Community program, Contract No. RII3-CT-2004-506078; the European Community-Research Infrastructure Integrating Activity "Study of Strongly Interacting Matter" (HadronPhysics2, Grant Agreement No. 227431) under the Seventh Framework Programme of EU; Austrian Federal Ministry of Science and Research BMBWK 650962/0001 VI/2/2009; Romanian National Authority for Scientific Research (ANCS); and the Grant-in-Aid for Specially Promoted Research (20002003), MEXT, Japan.

References

[1] S. Deser et al., Phys. Rev. 96 (1954) 774; T. L. Trueman, Nucl. Phys. 26 (1961) 57; A. Deloff, Phys. Rev. C 13 (1976) 730.
[2] B. Borasoy, R. Nißler, W. Weise, Phys. Rev. Lett. 94 (2005) 213401; Eur. Phys. J. A 25 (2005) 79.
[3] D. Jido, *et al.*, Nucl. Phys. A 725 (2003) 181.
[4] T. Hyodo and W. Weise, Phys. Rev. C 77 (2008) 035204.
[5] T. Hyodo, D. Jido, Prog. Part. Nucl. Phys.67 (2012) 55.
[6] Y. Akaishi, T. Yamazaki, Phys. Rev, C 65 (2002) 044005; T. Yamazaki, Y. Akaishi, Phys. Lett. B 535 (2002) 70; T. Yamazaki, Y. Akaishi, Phys. Rev. C 76 (2007) 045201.
[7] M. Agnello *et al.*, Phys. Rev. Lett. 94 (2005) 212303.
[8] T. Suzuki *et al.*, Phys. Rev. C 76 (2007) 068202.

[9] T. Yamazaki *et al.*, Phys. Rev. Lett. 104 (2010) 132502.
[10] C. Curceanu, J. Zmeskal, Mini-Proceedings of ECT Workshop “Strangeness in Nuclei”, arXiv:1104.1926 [nucl-ex] (2011), and references therein.
[11] J. D. Davies, *et al.*, Phys. Lett. B 83 (1979) 55.
[12] M. Izycki, *et al.*, Z. Phys. A 297 (1980) 11.
[13] P. M. Bird, *et al.*, Nucl. Phys. A 404 (1983) 482.
[14] M. Iwasaki, *et al.* Phys. Rev. Lett. 78 (1997) 3067; T. M. Ito, *et al.*, Phys. Rev. C 58 (1998) 2366.
[15] G. Beer, *et al.* (DEAR Collaboration), Phys. Rev. Lett. 94 (2005) 212302.
[16] W.E. Humphrey and R.R. Rose, Phys. Rev. 127 (1962) 1305.
[17] J.K. Kim, Phys. Rev. Lett. 19 (1967) 1074.
[18] A.D. Martin, Nucl. Phys. B179 (1981) 33.
[19] U.-G. Meißner, U. Raha, A. Rusetsky, Eur. Phys. J. C 35 (2004) 349.
[20] R. Nißler, PhD thesis, Univ. of Bonn (2007); B. Borasoy, U.-G. Meißner, R. Nißler, Phys. Rev. C 74(2006) 055201.
[21] J. A. Oller, J. Prades, M. Verbeni, Phys. Rev. Lett. 95 (2005) 172502.
[22] J. A. Oller, Eur. Phys. J. A 28 (2006) 63.
[23] B. Borasoy, R. Nißler, W. Weise, Phys. Rev. Lett. 96 (2006) 199201.
[24] J. A. Oller, J. Prades, M. Verbeni, Phys. Rev. Lett. 96 (2006) 199202.
[25] J. Révai, N. V. Shevchenko, Phys. Rev. C. 79 (2009) 035202.
[26] A. Cieplý, J. Smejkal, Eur. Phys. J. A 34 (2007) 237.
[27] N. V. Shevchenko, J. Révai, Few-Body Systems 44 (2008) 187.
[28] E. Oset, A. Ramos, Nucl. Phys. A 635 (1998) 99.
[29] W. Weise, Nucl. Phys. A 835 (2010) 51.
[30] J. Zmeskal, Prog. Part. Nucl. Phys. 61 (2008) 512.
[31] M. Bazzi, *et al.* (SIDDHARTA Collaboration), Phys. Lett. B 704 (2011) 113.
[32] C. Milardi *et al.*, Int. J. Mod. Phys. A, 24 (2009) 360.
[33] M. Zobov *et al.*, Phys. Rev. Lett., 104 (2010) 174801.
[34] T. Ishiwatari, *et al.*, Nucl. Instrum. Methods Phys. Res. A 556 (2006) 509.
[35] M. Bazzi, *et al.* (SIDDHARTA Collaboration), Phys. Lett. B 697 (2011) 199.
[36] M. Bazzi, *et al.* (SIDDHARTA Collaboration), Phys. Lett. B 681 (2009) 310.
[37] T. Koike, T. Harada, Y. Akaishi, Phys. Rev. C53 (1996) 79.
[38] A. Cieplý, E. Friedman, A. Gal, D. Gazda, J. Mareš, Phys. Rev. C 84 (2011) 045206.
[39] Y. Ikeda, T. Hyodo, W. Weise, Phys. Lett. B 706 (2011) 63.
[40] SIDDHARTA-2 Collaboration, Proposal of Laboratori Nazionali di Frascati of INFN, *The upgrade of the SIDDHARTA apparatus for an enriched scientific case* (2010).



Figure 6: A global simultaneous fit result of the X-ray energy spectra of hydrogen and deuterium data. (a) Residuals of the measured kaonic-hydrogen X-ray spectrum after subtraction of the fitted background, clearly displaying the kaonic-hydrogen K-series transitions. The fit components of the K^-p transitions are also shown, where the higher transitions, greater than $K\beta$, are summed. (b)(c) Measured energy spectra with the fit lines for each dataset. Fit components of the background X-ray lines and a continuous background are also shown. The dot-dashed vertical line indicates the EM value of the kaonic-hydrogen $K\alpha$ energy. (Note that the characteristic $K\alpha$ line consists of $K\alpha1$ and $K\alpha2$ lines, both of which are shown.)



Figure 7: Comparison of the present result for the strong-interaction 1*s*-energy-level shift and width of kaonic hydrogen with the two experimental results : KEK-PS E228 (1997) [14] and DEAR (2005) [15]. The error bars correspond to quadratically added statistical and systematic errors. The right panel shows the error in the energy shift as a function of the width (vertical axis) for each experiment. The dashed lines represent the SIDDHARTA precision calculated assuming the same statistics but with differing width.